arXiv:1011.1635v2 [math.AT] 3 Oct 2012

Kontsevich's swiss cheese conjecture

Justin D. Thomas

October 5, 2012

Abstract

We prove a conjecture of Kontsevich which states that if A is an E_{d-1} algebra then the Hochschild cohomology object of A is the universal E_d algebra acting on A. The notion of an E_d algebra acting on an E_{d-1} algebra was defined by Kontsevich using the swiss cheese operad of Voronov. The degree 0 and 1 pieces of the swiss cheese operad can be used to build a cofibrant model for A as an $E_{d-1} - A$ module. The theorem amounts to the fact that the swiss cheese operad is generated up to homotopy by its degree 0 and 1 pieces.

1 Introduction

In [Ger63] Gerstenhaber showed that the Hochschild cohomology $\mathrm{HH}^*(A)$ of an associative algebra A is a graded Lie algebra and a graded commutative algebra, and the two structures are compatible. Any chain complex with this algebraic structure is now called a Gerstenhaber algebra. In [Coh76] (see also [Sin06]) Fred Cohen showed that the operad controlling Gerstenhaber algebras is $H_{-*}(E_2)$. Deligne later asked if the action of $H_{-*}(E_2)$ on $\mathrm{HH}^*(A)$ descends from a natural action at the level of chains. In other words, is there a natural algebra structure of $\mathsf{Chains}(E_2)$ on $\mathrm{CH}^*(A)$ which recovers the structure discovered by Gerstenhaber after passing to (co)homology?

Already, this question is evidently in the realm of homotopy theory. So let us replace the associative algebras by E_1 algebras. This makes it clear that the question is fundamentally one about the relationship between the operads E_1 and E_2. Indeed, we can further consider the question in light of the relationship between E_d and E_{d-1} algebras. For any E_{d-1} algebra in a sufficiently rich homotopical category $\mathcal{C}$ we can make sense of its Hochschild cohomology. The Hochschild cohomology of A is denoted $\mathrm{Hoch}(A)$ and is an object of $\mathcal{C}$. This convention is a bit confusing, as it is the Hochschild *cochain complex* in the case where $\mathcal{C}$ is the category of differential graded vector spaces, but in a broader context it is best to just refer to it as the Hochschild cohomology of A.

The original Deligne conjecture where A is an E_1 algebra in the category of chain complexes has been solved several times [Tam98, BF04, MS02, KS00, Vor00] The generalized version where A is an E_d algebra in a general category like $\mathcal{C}$ has been proven by [HKV06, Lur11]. What we show here is that $\mathrm{Hoch}(A)$ is not just an E_d algebra, but comes equipped with a universal property. It is the universal E_d algebra acting on the E_{d-1} algebra A. The case $d = 2$ in the algebraic setting can be found in [DTT09].

The notion of an E_d algebra acting on an E_{d-1} algebra was introduced in [Kon99]. This notion uses the swiss cheese operad SC_d of Voronov [Vor99]. This is a two-colored operad which interpolates between E_d and E_{d-1}. A swiss cheese algebra is a pair (B, A) where B is an E_d algebra, A is an E_{d-1} algebra, and there is some extra structure compatible with these (definition 13). We refer to this extra structure as an *action* of B on A.

The case $d = 1$ is enlightening. For simplicity, let us work in the category of vector spaces. A (non-unital) E_0 algebra A in vector spaces is just a vector space with no extra data. The Hochschild cohomology in this case is $\hom(A, A)$. If B is an associative algebra, it is in particular an E_1 algebra. An SC_1 structure on the (B, A) pair (B, A) then amounts to the choice of a B-module structure on A.

In this case, the swiss cheese conjecture merely states that $\hom(A, A)$ is an associative algebra, and giving an SC_1 structure on (B, A) is equivalent to giving a map of associative algebras $B \to \mathrm{Hoch}(A) = \hom(A, A)$. We prove the analog of this when B is an E_d algebra and A is an E_{d-1} algebra, $d \geq 1$.

1.1 Outline of the paper

In section 2 we define the E_d and SC_d operads, and give both an imprecise and a precise statement of the theorem we will prove. We also outline the idea of the proof. In section 3 we define Hochschild cohomology for E_{d-1} algebras and show that we can use the swiss cheese operad to construct a model for Hochschild cohomology. We use this model in section 4 to prove a "universal cheese" theorem which applies to an arbitrary operad acting on Hochschild cohomology. Finally, in section 5 we show that E_d does indeed act, up to homotopy, on Hochschild cohomology, and the "universal cheese" theorem specializes to the swiss cheese theorem.

2 The swiss cheese operad

Fix a set K, a K-colored set is a pair (I, col) where I is a set and $\mathrm{col} : I \to K$ is a map of sets, called the *coloring*. We will often denote such a colored set simply by I, leaving the coloring implicit. Let $\mathrm{Aut}(I)$ be the group of bijections on the set I which preserve its coloring. The following definition follows [BV73].

Definition 1. Let $(\mathcal{S}, \otimes)$ be a symmetric monoidal category containing all coproducts and assume that $\otimes$ distributes over coproducts. Let K be a set. A K-colored operad $\mathcal{O}$ in $\mathcal{S}$ is a symmetric monoidal category enriched over $\mathcal{S}$ whose objects are K-colored finite sets, $I \to K$. We require that, on objects, the symmetric monoidal structure of $\mathcal{O}$ is the disjoint union of sets over K. This endows each hom object $\mathcal{O}(I; J)$ with the structure of a right $\mathrm{Aut}(I)$ module and a left $\mathrm{Aut}(J)$ module. We also require each of the canonical maps

$$\coprod_{f\colon I \to J} \bigotimes_{j \in J} \mathcal{O}(f^{-1}(j); \{j\}) \to \mathcal{O}(I; J), \tag{2}$$

to be isomorphisms of right $\mathrm{Aut}(I)$ and left $\mathrm{Aut}(J)$ modules. The maps $f\colon I \to J$ need not preserve the colorings. Note that the input finite set is separated from the output finite set by a semicolon. This is to reduce confusion in forthcoming notation.

Notation 3. Let $\underline{n}$ to denote the finite set $\{1, \ldots, n\}$. Typically K will be $K = \{\mathrm{f}, \mathrm{h}\}$, where f stands for *full disc* and h stands for *half disc.* In this case we use (n, m) to denote the K-colored set which is the disjoint union of

$$\underline{n} \to \{\mathrm{f}\} \text{ and } \underline{m} \to \{\mathrm{h}\}.$$

If $K \simeq \{*\}$, then a K-colored operad will simply be called a 1-colored operad. Any 1-colored operad $\mathcal{E}$ gives for each $m, m' \in \mathbb{Z}_{\geq 0}$, objects $\mathcal{E}(\underline{m}'; \underline{m}) \in \mathcal{S}$. We denote $\mathcal{E}(\underline{m}'; \underline{m})$ simply by $\mathcal{E}(m'; m)$ and $\mathcal{E}(m; 1)$ simply by $\mathcal{E}(m)$.

Any $\{\mathrm{f}, \mathrm{h}\}$-colored operad $\mathcal{O}$ gives for each $n, m, n', m' \in \mathbb{Z}_{\geq 0}$, objects

$$\mathcal{O}(n', m'; n, m) \coloneqq \mathcal{O}((n', m'); (n, m)) \in \mathcal{S}.$$

We denote $\mathcal{O}(n, m; 0, 1)$ by $\mathcal{O}^{\mathrm{h}}(n, m)$ and we denote $\mathcal{O}(n, m; 1, 0)$ by $\mathcal{O}^{\mathrm{f}}(n, m)$.

Example 4. We spell out definition 1 in the 1-colored case. In a 1-colored operad $\mathcal{E}$, the right $\mathrm{Aut}(\underline{m})$ modules $\mathcal{E}(m)$, $m \geq 0$, determine the objects $\mathcal{E}(m'; m)$ by formula 2 in definition 1. The categorical composition laws (i.e. operadic structure) of $\mathcal{E}$ are $\mathcal{S}$-morphisms

$$\mathcal{E}(m'; m'') \otimes \mathcal{E}(m; m') \to \mathcal{E}(m; m''),$$

where $m, m', m'' \in \mathbb{Z}_{\geq 0}$. These maps make $\mathcal{E}$ into a symmetric monoidal category enriched over $\mathcal{S}$. It is enough to give the maps where $m'' = 1$.

$$\mathcal{E}(m') \otimes \coprod_{m = k_1 + \cdots + k_{m'}} \mathcal{E}(k_1) \otimes \cdots \otimes \mathcal{E}(k_{m'}) \to \mathcal{E}(m).$$

The equivariance and associativity conditions [May72, definition 1.1] [MSS07, definition 1.4] on the above maps are packaged in the requirement that $\mathcal{E}$ is a symmetric monoidal category.

Example 5. Let K be the one-point set $\{\mathrm{f}\}$ and let $(\mathcal{S}, \otimes)$ be the symmetric monoidal category $(\mathsf{Top}, \times)$ of compactly generated spaces with the cartesian product. The operad E_d is an $\{\mathrm{f}\}$-colored operad in the category Top. Let $\bar{D}^d$ be the closed unit disc inside $\mathbb{R}^d$. Call a map $f : \bar{D}^d \to \bar{D}^d$ a *little full-disc* (or *little d disc* or simply *little disc*) if f is of the form $f(x) = rx + c$ for some $0 < r \leq 1$ and $c \in \mathbb{R}^d$. Given a finite set I, the underlying set of $E_d(I; \mathrm{f})$ is the set of embeddings

$$f \colon \coprod_{i \in I} \bar{D}^d \to \bar{D}^d,$$

where each restriction $f_i \colon \bar{D}^d \to \bar{D}^d$ is a little full-disc. Using notation 3, any isomorphism $I \to \underline{n}$ induces an isomorphism $E_d(I; \mathrm{f}) \simeq E_d(n)$, and the latter can naturally be considered as a subset of $\mathbb{R}^{n+dn}$. This gives each $E_d(I; \mathrm{f})$ a topology. The operadic structure is given by composing little d discs as maps $\bar{D}^d \to \bar{D}^d$. The identity of E_d is the little full-disc $\mathsf{id} \colon \bar{D}^d \to \bar{D}^d$. This is the unital version of E_d, so $E_d(0) = *$ and $E_d(1)$ consists of more than just the identity.

Example 6. Let $K = \{\mathrm{f}, \mathrm{h}\}$. The K-colored operad SC_d is called the (d-dimensional) swiss cheese operad and is the principal subject of this paper. Like example 5 it is an operad in $(\mathsf{Top}, \times)$.

Let $\{\mathrm{f}\}$ and $\{\mathrm{h}\}$ denote the evident singleton K-colored sets. By definition 1, formula 2, we only need to define the spaces $SC_d(I; \mathrm{f})$ and $SC_d(I; \mathrm{h})$ for every K-colored set $(I, \mathrm{col} \colon I \to K)$. First, we define the "full-disc output" part of SC_d,

$$SC_d(I; \mathrm{f}) = \begin{cases} E_d(I; \mathrm{f}) & \mathrm{col}^{-1}(\mathrm{f}) = I \\ \emptyset & \text{else.} \end{cases}$$

To define the "half-disc output" part of SC_d, that is $SC_d(I; \mathrm{h})$, we first need the notion of *little half-discs*. Let $\bar{D}^d_+$ be the closed d-dimensional half-disc,

$$\bar{D}^d_+ = \{(x_1, \ldots, x_d) \in \mathbb{R}^d \mid |x| \leq 1 \text{ and } x_d \geq 0\}.$$

A *little half-disc* is defined to be a map $f \colon \bar{D}^d_+ \to \bar{D}^d_+$ of the form $f(x) = rx + c$ for some $0 < r \leq 1$ and $c \in \mathbb{R}^{d-1} \times \{0\}$. As a set, we define $SC_d(I; \mathrm{h})$ to consist of embeddings

$$f \colon \coprod_{i \in I} \bar{D}^d_i \to \bar{D}^d_+$$

where $\bar{D}^d_i$ is the full disc $\bar{D}^d$ if $\mathrm{col}(i) = \mathrm{f}$ and $\bar{D}^d_i$ is the half-disc $\bar{D}^d_+$ if $\mathrm{col}(i) = \mathrm{h}$. For each $i \in I$ the restriction $f_i \colon \bar{D}^d_i \to \bar{D}^d_+$ of the embedding f is required to be either a little full-disc (example 5) or little half-disc, depending on what $\bar{D}^d_i$ is. It is clear that if $\left|\mathrm{col}^{-1}(\mathrm{f})\right| = n$ and $\left|\mathrm{col}^{-1}(\mathrm{h})\right| = m$, then $SC_d(I; \mathrm{h})$ can be naturally embedded inside $\mathbb{R}^N$

where $N = n + dn + m + (d-1)m$. We give $SC_d(I; \mathrm{h})$ the subspace topology inherited from such an embedding.

Following notation 3, a point in $SC_d^{\mathrm{h}}(n,m)$ is given by n labeled full-discs and m labeled half-discs in the unit half-disc where none of the discs intersect and the half-discs all lie on the bottom. We allow the degenerate configuration when $(n,m) = (0,1)$ which is the unit half-disc contained in itself. Note that we have $SC_d^{\mathrm{h}}(0,0) = *$ and $SC_d^{\mathrm{h}}(1,0)$ contains more than one point. Thus we are using the unital swiss cheese operad. This differs from Kontsevich in [Kon99] and Voronov in [Vor99].

Composition in SC_d is given by substituting full-discs and half-discs into each other as in figure 6. More precisely, we have maps

$$E_d(n) \times E_d(k_1) \times \cdots \times E_d(k_n) \to E_d(k_1 + \cdots + k_n) \tag{7}$$

and

$$\begin{aligned} SC_d^{\mathrm{h}}(n,m) \times E_d(k_1) \times \cdots \times E_d(k_n) \times SC_d^{\mathrm{h}}(k_{n+1}, \ell_1) \times \cdots \times SC_d^{\mathrm{h}}(k_{n+m}, \ell_m) \\ \longrightarrow SC_d^{\mathrm{h}}(k_1 + \cdots + k_{n+m}, \ell_1 + \cdots + \ell_m). \end{aligned} \tag{8}$$



$$SC_d^{\mathrm{h}}(2,2) \times E_d(2) \times E_d(3) \times SC_d^{\mathrm{h}}(1,1) \times SC_d^{\mathrm{h}}(0,2) \to SC_d^{\mathrm{h}}(6,3)$$

Figure 1: The operadic composition for SC_d when $d = 2$. Full-discs or half-discs can take inputs. The full-discs and half-discs should be labelled independently. We have omitted labels in the figure.

Notice that we can identify $SC_d^{\mathrm{h}}(0,m)$ with $E_{d-1}(m)$ so that the restriction of SC_d to the spaces $SC_d^{\mathrm{h}}(0,\bullet)$ is isomorphic to the operad E_{d-1}. We say that E_{d-1} is the h color of SC_d and E_d is the f color of SC_d. We think of SC_d as interpolating between E_d and E_{d-1}.

Definition 9. Suppose $\mathcal{O}$ is K-colored operad in $\mathcal{S}$ and $\mathcal{C}$ is a symmetric monoidal category enriched over $\mathcal{S}$. An *algebra* over $\mathcal{O}$ in the category $\mathcal{C}$ is a symmetric monoidal functor $\mathcal{O} \to \mathcal{C}$. A morphism of $\mathcal{O}$ algebras is a monoidal natural transformation. The category of $\mathcal{O}$ algebras in $\mathcal{C}$ will be denoted $\mathrm{Alg}_{\mathcal{O}}(\mathcal{C})$.

Example 10. If $\mathcal{C}$ is a symmetric monoidal category enriched over Top, we can consider algebras over SC_d in $\mathcal{C}$. Such an algebra gives the data of a pair (B, A) of objects in $\mathcal{C}$ together with maps of topological spaces

$$E_d(n) \to \mathrm{map}_{\mathcal{C}}(B^{\otimes n}, B)$$

and

$$SC_d^{\mathrm{h}}(n,m) \to \mathrm{map}_{\mathcal{C}}(B^{\otimes n} \otimes A^{\otimes m}, A),$$

where $\mathrm{map}_{\mathcal{C}}(C, C')$ is the topological space of maps between two objects C, C' in $\mathcal{C}$. These data must satisfy conditions guaranteeing they assemble into a symmetric monoidal functor $SC_d \to \mathcal{C}$.

The object B corresponds to the object $\{\mathrm{f}\}$ of SC_d and the object A corresponds to $\{\mathrm{h}\}$. Together these form a K-colored operad $\mathrm{End}(B, A)$ in Top where, using notation 3,

$$\mathrm{End}^{\mathrm{f}}(B, A)(n, m) = \mathrm{map}(B^{\otimes n} \otimes A^{\otimes m}, B) \tag{11}$$

$$\mathrm{End}^{\mathrm{h}}(B, A)(n, m) = \mathrm{map}(B^{\otimes n} \otimes A^{\otimes m}, A). \tag{12}$$

The collection of SC_d algebra structures on a fixed pair (B, A) as the collection of symmetric monoidal functors

$$SC_d \to \mathrm{End}(B, A),$$

which restrict to the identity on the set of objects. Simply put, a swiss cheese algebra (B, A) is an E_d algebra B, an E_{d-1} algebra A, and some chosen mixing of these structures. We refer to this mixing as an action of B on A. The following definition is due to Kontsevich [Kon99].

Definition 13. Let B be an E_d algebra and A an E_{d-1} algebra. A *swiss cheese action of B on A* is the structure of a swiss cheese algebra on the pair (B, A) extending the given E_d and E_{d-1} structures. We may also simply call this "an action of B on A".

With this we can informally state the conjecture proven in this paper.

Theorem (Informal statement of Kontsevich's Swiss Cheese Conjecture)**.** The Hochschild cohomology $\mathrm{Hoch}(A)$ of an E_{d-1} algebra A is the universal E_d algebra acting on A. In other words, for any E_{d-1} algebra A, there is an E_d algebra structure on $\mathrm{Hoch}(A)$ such that for any E_d algebra B, giving a map of E_d algebras $B \to \mathrm{Hoch}(A)$ is equivalent to giving the structure of an SC_d algebra on the pair (B, A) extending the given E_d and E_{d-1} structures.

The basic structure of the proof is outlined in diagram 14. The categories of operads shown in the diagram are defined precisely in section 2.1. Informally, $\mathsf{Op}(\mathrm{Coll})$ consists of K-colored operads whose f-colored output is trivial, and $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$ further restricts to those pieces whose f-colored inputs total 0 or 1. The forgetful functors are presented below as straight arrows; there is a left adjoint shown as a bent arrow.

Boardman and Vogt's W construction [BV73], [BM06] is an explicit cofibrant replacement functor, which we apply to SC_d, to get an equivalent cofibrant operad $\mathsf{SC_d} \coloneqq WSC_d$. The W construction does not strictly commute with the forgetful functors in 14. In particular $W(SC_d^{\mathrm{h}})$ is not isomorphic to $(WSC_d)^{\mathrm{h}}$, but they are homotopy equivalent. We will let $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ denote $W(SC_d^{\mathrm{h}})$ and $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}} = W(SC_d^{\mathrm{h1}})$.

$$\begin{tikzcd}
 & \mathsf{Op}_K \arrow[r] & \mathsf{Op}(\mathrm{Coll}) \arrow[r] & \mathsf{Op}(\mathrm{Coll}^{\leq 1}) \arrow[l, bend right] \\
 & \mathsf{SC_d} \arrow[r, mapsto] & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}} \arrow[r, mapsto] & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}} \arrow[dl, mapsto] \\
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes \mathsf{E} & & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \arrow[ll, dashed, "\text{transfer of}", "\text{structure}"'] &
\end{tikzcd} \tag{14}$$

Proposition 53 shows that we can use $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ to construct a model for Hochschild cohomology. This allows us to prove a weak version of the swiss cheese theorem in proposition 55 taking place in the context of $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$. Next, we will take the free extension of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}} \in \mathsf{Op}(\mathrm{Coll}^{\leq 1})$ to an operad in $\mathsf{Op}(\mathrm{Coll})$, to get $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$. This immediately gives a version of the swiss cheese theorem in the context of $\mathsf{Op}(\mathrm{Coll})$, see corollary 62. Then we use the fact that $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ is freely generated by its degree 0 and 1 pieces to prove a version of the swiss cheese theorem in the context of Op_K. None of these three versions of the swiss cheese theorem make any use of E_d. One can think of this last "universal cheese" theorem (proposition 75) as a construction of the universal K-colored operad constructed from $\mathsf{E_{d-1}}$ and controlling $\mathsf{E_{d-1}}$-linear actions on A.

To bring E_d back in to the story, we use a technical result, theorem 77, which shows that the canonical map $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ is an acyclic cofibration. Observe that one can view $\mathsf{SC_d}$ as

$\mathsf{SC}_\mathsf{d}^\mathrm{h}$ equipped with the extra structure of a homotopy right action of E_d. Now use a transfer of structure argument to construct an operad E which is equivalent to E_d and which acts on the right on $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$. This allows us to define $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$, which we show is equivalent to SC_d in 85. The universal property of $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$ with respect to Hochschild cohomology is stated in theorem 15 and follows from proposition 75.

Theorem 15 (Precise version of Kontsevich's Swiss Cheese Conjecture)**.** Let $\mathcal{C}$ be a symmetric monoidal model category tensored over Top and satisfying the conditions in notation 33. Let $A \in \mathrm{Alg}_{\mathsf{E}_{\mathsf{d}-1}}(\mathcal{C})$ be cofibrant and fibrant using the projective model structure (32). There is a model of SC_d, called $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$ where $\mathsf{E} \simeq E_d$. There is also a model for the Hochschild cohomology object of A, called $\mathrm{Hoch}(A)$, such that $\mathrm{Hoch}(A)$ is the universal E algebra acting on A through $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$. That is, $\mathrm{Hoch}(A)$ is an E algebra and this structure, together with the E_{d-1} algebra structure on A, can be extended to an $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$ algebra structure on $(\mathrm{Hoch}(A), A)$ in such a way that there is an isomorphism of categories

$$\mathrm{Alg}^{A}_{(\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E})}(\mathcal{C}) \cong \mathrm{Alg}_{\mathsf{E}}(\mathcal{C})_{/\mathrm{Hoch}(A)}.$$

The category on the left consists of E algebras B together with an action of B on A. The category on the right consists of E algebras B together with an E algebra map $B \to \mathrm{Hoch}(A)$.

2.1 Defining SC_d^h and $SC_d^{\mathrm{h}1}$

Recall that $\mathcal{C}$ is a symmetric monoidal category enriched over $\mathcal{S}$, our basic category in which our operads live. We will assume that both $\mathcal{C}$ and $\mathcal{S}$ have all coproducts and that tensor products distribute over coproducts. In the case of operads from Op_K we have $(\mathcal{S}, \otimes) = (\mathsf{Top}, \times)$, the symmetric monoidal category of compactly generated topological spaces with cartesian product. In the case of operads such as SC_d^h and $\mathsf{SC}_\mathsf{d}^\mathrm{h}$ from $\mathsf{Op}(\mathrm{Coll})$, we have $(\mathcal{S}, \otimes) = (\mathrm{Coll}(\mathsf{Top}), \otimes)$ from definition 16. Finally, for $\mathsf{SC}_\mathsf{d}^{\mathrm{h}1}$ and $SC_d^{\mathrm{h}1}$ we use $\mathcal{S} = \mathrm{Coll}^{\leq 1}(\mathsf{Top})$ as in definition 17.

Definition 16. Let Σ denote the category of finite sets with morphisms given by bijections. The category of functors $\Sigma^{op} \to \mathcal{C}$, denoted $\mathrm{Coll}(\mathcal{C})$, is usually called the category of *collections* in $\mathcal{C}$. We endow $\mathrm{Coll}(\mathcal{C})$ with the usual symmetric monoidal structure given by left Kan extension of $\Sigma^{op} \times \Sigma^{op} \to \mathcal{C} \times \mathcal{C} \to \mathcal{C}$ along disjoint union of sets $\Sigma^{op} \times \Sigma^{op} \to \Sigma^{op}$. Specifically, if $X, Y \in \mathrm{Coll}(\mathcal{C})$, then $(X \otimes Y) \in \mathrm{Coll}(\mathcal{C})$ satisfies

$$(X \otimes Y)(\underline{n}) = \coprod_{n = n_1 + n_2} \mathrm{Ind}^{\mathrm{Aut}(\underline{n})}_{\mathrm{Aut}(\underline{n}_1) \times \mathrm{Aut}(\underline{n}_2)} (X(\underline{n}_1) \otimes_\mathcal{C} Y(\underline{n}_2)),$$

where $\otimes$ is the tensor product in $\mathrm{Coll}(\mathcal{C})$, and $\otimes_\mathcal{C}$ is the tensor product in $\mathcal{C}$. See [Har07, definition 3.3] for more details. We will abbreviate $\mathrm{Coll}(\mathsf{Top})$ as Coll.

Definition 17. Let $\Sigma^{\leq 1} \subset \Sigma$ denote the full subcategory of finite sets of size 0 or 1 together with bijections as morphisms. Let $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ denote the symmetric monoidal category of functors $(\Sigma^{\leq 1})^{op} \to \mathcal{C}$, with monoidal structure inherited from $\mathrm{Coll}(\mathcal{C})$. Call these the *degree 0-1 collections in* $\mathcal{C}$. Concretely, $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ is just the category $\mathcal{C} \times \mathcal{C}$ endowed with the symmetric monoidal structure

$$(C_0, C_1) \otimes (D_0, D_1) = (C_0 \otimes D_0, (C_0 \otimes D_1) \coprod (C_1 \otimes D_0)). \tag{18}$$

The braiding isomorphism $(C_0, C_1) \otimes (D_0, D_1) \to (D_0, D_1) \otimes (C_0, C_1)$ is induced from the braiding isomorphism on $\mathcal{C}$. We will abbreviate $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$ as $\mathrm{Coll}^{\leq 1}$.

Definition 19. Let Op_K denote the category of $K = \{\mathrm{f}, \mathrm{h}\}$-colored operads in Top. There is a forgetful functor $\mathsf{Op}_K \to \mathsf{Op}(\mathrm{Coll})$. This functor takes $\mathcal{O} \in \mathsf{Op}_K$ to the operad $\mathcal{O}^\mathrm{h}$ whose arity m component is the collection $n \mapsto \mathcal{O}^\mathrm{h}(n, m)$ (see notation 3). We forget the spaces $\mathcal{O}^e(n, m)$, and think of elements of $\mathcal{O}^h(n, m)$ as degree n, arity m elements of $\mathcal{O}^h$.

The functor $\mathsf{Op}(\mathrm{Coll}) \to \mathsf{Op}(\mathrm{Coll}^{\leq 1})$ is induced by the symmetric monoidal functor $\mathrm{Coll} \to \mathrm{Coll}^{\leq 1}$. Denote the image of $\mathcal{O}$ in $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$ as $\mathcal{O}^{\mathrm{h}1}$.

Example 20. We outline the structure of $\mathsf{SC}_\mathsf{d}^\mathrm{h}$ as an operad from $\mathsf{Op}(\mathrm{Coll})$. Think of $\mathsf{SC}_\mathsf{d}^\mathrm{h}(\bullet, m)$ as the collection $n \mapsto \mathsf{SC}_\mathsf{d}^\mathrm{h}(n, m)$. In 21, $\otimes$ is the tensor product of collections. The operad composition law is

$$\mathsf{SC}_\mathsf{d}^\mathrm{h}(\bullet, m) \otimes \mathsf{SC}_\mathsf{d}^\mathrm{h}(\bullet, \ell_1) \otimes \cdots \otimes \mathsf{SC}_\mathsf{d}^\mathrm{h}(\bullet, \ell_m) \to \mathsf{SC}_\mathsf{d}^\mathrm{h}(\bullet, \ell), \tag{21}$$

where $\ell = \sum_i \ell_i$. The degree n component of the right hand collection is $\mathsf{SC}_\mathsf{d}^\mathrm{h}(n, \ell)$. The degree n component of the left hand collection is

$$\coprod_{n_0+\cdots+n_m=n} \mathrm{Ind}\ \mathsf{SC}_\mathsf{d}^\mathrm{h}(n_0, m) \times \mathsf{SC}_\mathsf{d}^\mathrm{h}(n_1, \ell_1) \times \cdots \times \mathsf{SC}_\mathsf{d}^\mathrm{h}(n_m, \ell_m),$$

where Ind is the induction functor giving the correct symmetric group action. The point is that if we delete all appearances of E_d from 8 then it provides exactly the data of 21.

Example 22. We outline the structure of $SC_d^{\mathrm{h}1}$ as a 1-colored operad from $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$. For each m', m, define an object $SC_d^{\mathrm{h}1}(m'; m) \in \mathrm{Coll}^{\leq 1}(\mathcal{C})$,

$$\begin{aligned} SC_d^{\mathrm{h}1}(m'; m) &= (SC_d^{\mathrm{h}}(0, m'; 0, m), SC_d^{\mathrm{h}}(1, m'; 0, m)) \qquad (23)\\ &\cong (E_{d-1}(m'; m), SC_d^{\mathrm{h}}(1, m'; 0, m)). \qquad (24)\end{aligned}$$

Using the symmetric monoidal structure from definition 17 and the identification in equation 24 we can write the operad structure maps on $SC_d^{\mathrm{h}1}$ as a triple of morphisms. The map of degree 0 pieces,

$$E_{d-1}(m''; m) \times E_{d-1}(m'; m'') \to E_{d-1}(m'; m),$$

and the maps of degree 1 pieces,

$$E_{d-1}(m''; m) \times SC_d^{\mathrm{h}}(1, m'; 0, m'') \to SC_d^{\mathrm{h}}(1, m'; 0, m),$$

and

$$SC_d^{\mathrm{h}}(1, m''; 0, m) \times E_{d-1}(m'; m'') \to SC_d^{\mathrm{h}}(1, m'; 0, m).$$

3 Hochschild cohomology from swiss cheese

For the remainder of the paper we replace E_{d-1}, E_d, and SC_d by cofibrant models given by the Boardman-Vogt W construction [BV73], [BM06]. We will denote these cofibrant replacements by $\mathsf{E}_{\mathsf{d-1}}, \mathsf{E}_\mathsf{d}$, and SC_d. We also want to restrict our attention to swiss cheese algebras in categories where we can do homotopy theory. In the proper context (notation 33), the Hochschild cohomology of an $\mathsf{E}_{\mathsf{d-1}}$ algebra A has a natural model constructed from A and the degree 0-1 parts of $\mathsf{SC}_\mathsf{d}^\mathrm{h}$ (proposition 53).

3.1 Homotopy theoretic context

Definition 25. From [Hov99, definition 4.2.6], a *symmetric monoidal model category* $\mathcal{S}$ is a closed symmetric monoidal category whose monoidal structure $\otimes\colon \mathcal{S} \times \mathcal{S} \to \mathcal{S}$ is a Quillen bifunctor. We assume the monoidal unit of $\mathcal{S}$ is cofibrant.

Example 26. The category $(\mathsf{Top}, \times)$ of compactly generated spaces with the cartesian product and Serre model structure is a symmetric monoidal model category.

Definition 27. Let $\mathcal{S}$ be a symmetric monoidal model category. A *symmetric monoidal model category tensored over* $\mathcal{S}$ is a closed symmetric monoidal model category $\mathcal{C}$, together with a symmetric monoidal Quillen functor $\mathcal{S} \to \mathcal{C}$ (See [Hov99, definition 4.2.20]).

In particular, $\mathcal{C}$ comes equipped with functors

$$\otimes : \mathcal{C} \times \mathcal{C} \to \mathcal{C} \quad \otimes : \mathcal{S} \times \mathcal{C} \to \mathcal{C}.$$
$$\underline{\hom}_{\mathcal{C}} : \mathcal{C}^{op} \times \mathcal{C} \to \mathcal{C} \quad \operatorname{map}_{\mathcal{C}} : \mathcal{C}^{op} \times \mathcal{C} \to \mathcal{S}$$

The mapping spaces $\operatorname{map}_{\mathcal{C}}(A, B)$ give $\mathcal{C}$ the structure of a category enriched over $\mathcal{S}$, so we can speak of $\mathsf{E_{d-1}}, \mathsf{E_d}$ and $\mathsf{SC_d}$ algebras in $\mathcal{C}$.

For any object A of $\mathcal{C}$, the functor $- \otimes A$ has right adjoints $\underline{\hom}_{\mathcal{C}}(A, -) : \mathcal{C} \to \mathcal{C}$ and $\operatorname{map}_{\mathcal{C}}(A, -) : \mathcal{C} \to \mathcal{S}$. This data satisfies Quillen's SM7 axiom ([Hov99, section 4.2]). We assume that the unit of $\mathcal{C}$ is cofibrant.

Example 28. The category $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ from definition 17 is tensored over the symmetric monoidal category $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$ with

$$\mathrm{Coll}^{\leq 1}(\mathsf{Top}) \otimes \mathrm{Coll}^{\leq 1}(\mathcal{C}) \to \mathrm{Coll}^{\leq 1}(\mathcal{C})$$

given by the analogue of equation 18,

$$(X_0, X_1) \otimes (C_0, C_1) = (X_0 \otimes C_0, X_0 \otimes C_1 \coprod X_1 \otimes C_0),$$

where $X_i \in \mathsf{Top}$ and $C_i \in \mathcal{C}$.

Example 29. The symmetric monoidal functor $\mathsf{Top} \to \mathrm{Coll}^{\leq 1}(\mathsf{Top})$ sending X to $(X, \emptyset)$ makes both $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$ and $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ into symmetric monoidal model categories tensored over Top. Note that for a topological 1-colored operad $\mathcal{O}$, we can consider algebras over $\mathcal{O}$ in $\mathcal{C}$ as well as algebras over $\mathcal{O}$ in $\mathrm{Coll}^{\leq 1}(\mathcal{C})$.

The category of degree 0-1 collections is naturally home to $\mathcal{O}$-algebras A and $\mathcal{O}$-A modules M.

Definition 30. Suppose $\mathcal{O}$ is a 1-colored operad in Top, and let (A, M) be an object of $\mathrm{Coll}^{\leq 1}(\mathcal{C})$. The structure of an $\mathcal{O}$ algebra on the degree 0-1 collection (A, M) is the structure of an $\mathcal{O}$ algebra on A together with the data of maps

$$\mathcal{O}(m) \otimes M \otimes A^{\otimes m-1} \to M,$$

satisfying certain conditions. We call this data the structure of an $\mathcal{O}$-A *module* on M. Given a fixed $\mathcal{O}$ algebra $A \in \mathcal{C}$, the category $\mathrm{Mod}^A_{\mathcal{O}}(\mathcal{C})$ of $\mathcal{O}$-A modules has objects $M \in \mathcal{C}$ together with the structure of an $\mathcal{O}$ algebra on $(A, M) \in \mathrm{Coll}^{\leq 1}(\mathcal{C})$ extending the given $\mathcal{O}$ algebra structure on A.

We can enrich $\mathrm{Mod}^A_{\mathcal{O}}(\mathcal{C})$ over $\mathcal{C}$ and over Top. Indeed, given $\mathcal{O}$-A modules M' and M, we can define the hom-object of $\mathcal{O}$-A module morphisms from M' to M as the equalizer

$$\underline{\hom}^A_{\mathcal{O}}(M', M) \xrightarrow{\text{eq}} \underline{\hom}_{\mathcal{C}}(M', M) \rightrightarrows \underline{\hom}_{\mathcal{C}}(F^A_{\mathcal{O}}(M'), M).$$

Here $F^A_{\mathcal{O}} : \mathcal{C} \to \mathcal{C}$ is the free $\mathcal{O}$-A module monad on $\mathcal{C}$. One can define thus using the monad $F_{\mathcal{O}}$ on $\mathcal{C} \times \mathcal{C}$ which sends (A, M) to $(F^0_{\mathcal{O}}(A), F^1_{\mathcal{O}}(A, M))$, where $F^0_{\mathcal{O}}$ is the free $\mathcal{O}$-algebra monad and

$$F^1_{\mathcal{O}}(A, M) = \coprod_m \mathcal{O}(m) \otimes_{S_m} \Big(\coprod_{i=1}^m A^{\otimes i-1} \otimes M \otimes A^{\otimes m-i} \Big).$$

The natural transformation $F^1_{\mathcal{O}}(F^0_{\mathcal{O}}(-), F^1_{\mathcal{O}}(-,-)) \to F^1_{\mathcal{O}}(-,-)$ is given by composition in $\mathcal{O}$. This, along with $F^0_{\mathcal{O}} F^0_{\mathcal{O}} \to F^0_{\mathcal{O}}$, defines the monad structure $F_{\mathcal{O}} F_{\mathcal{O}} \to F_{\mathcal{O}}$. Finally, we define the free $\mathcal{O}$-A algebra monad via the coequalizer

$$F^1_{\mathcal{O}}(F^0_{\mathcal{O}}(A), F^1_{\mathcal{O}}(A, M)) \rightrightarrows F^1_{\mathcal{O}}(A, M) \xrightarrow{\text{coeq}} F^A_{\mathcal{O}}(M).$$

It is clear that every $\mathcal{O}$-A module M is equipped with a canonical map $F^A_{\mathcal{O}}(M) \to M$. The two parallel arrows in the equalizer are given by the two maps $F^A_{\mathcal{O}}(M') \to M'$ and $F^A_{\mathcal{O}}(M) \to M$. The topological space $\operatorname{map}^A_{\mathcal{O}}(M', M)$ is defined as an equalizer in exactly the same manner.

Example 31. Let $A \in \mathcal{C}$ be an $\mathsf{E_{d-1}}$ algebra, then the degree 0-1 collection (A, A) is naturally an $\mathsf{E_{d-1}}$ algebra. That is, A is naturally an $\mathsf{E_{d-1}}$-A module.

Suppose we have an adjunction $F\colon \mathcal{D} \leftrightarrows \mathcal{E}\colon U$ where F is the left adjoint; U is the right adjoint; $\mathcal{D}$ is a cofibrantly generated model category; and $\mathcal{E}$ has enough limits and colimits. There are conditions ([BM03, section 2.5]) on such an adjunction guaranteeing that $\mathcal{E}$ inherits a model structure where f in $\mathcal{E}$ is weak equivalence or fibration if and only if $G(f)$ in $\mathcal{D}$ is a weak equivalence or fibration, respectively. We are not concerned with these conditions here, but we assume that they apply to the adjunctions listed in definition 32.

Definition 32. Recall from example 22 that SC_d^{h1} is a 1-colored operad in $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$. Let $\mathsf{SC_d^{h1}} \in \mathrm{Coll}^{\leq 1}(\mathsf{Top})$ be $W(SC_d^{\mathrm{h1}})$. Since $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ is enriched over $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$ by example 28, we can consider $\mathsf{SC_d^{h1}}$ algebras in $\mathrm{Coll}^{\leq 1}(\mathcal{C})$. In addition, by example 29 we can consider $\mathsf{E_{d-1}}$ algebras in $\mathrm{Coll}^{\leq 1}(\mathcal{C})$. There are adjunctions

$$\mathcal{C} \leftrightarrows \mathrm{Alg}_{\mathsf{E_{d-1}}}(\mathcal{C}) \quad \mathcal{C} \leftrightarrows \mathrm{Mod}^{A}_{\mathsf{E_{d-1}}}(\mathcal{C})$$
$$\mathcal{C} \times \mathcal{C} \leftrightarrows \mathrm{Alg}_{\mathsf{E_{d-1}}}(\mathrm{Coll}^{\leq 1}(\mathcal{C})) \leftrightarrows \mathrm{Alg}_{\mathsf{SC_d^{h1}}}(\mathrm{Coll}^{\leq 1}(\mathcal{C}))$$

We describe the right adjoints only. On the top left the $\mathsf{E_{d-1}}$ algebra A is sent to the underlying object A of $\mathcal{C}$. The top right functor sends the $\mathsf{E_{d-1}}$-A module M to the underlying object M of $\mathcal{C}$. In the pair of composeable adjunctions, an $\mathsf{SC_d^{h1}}$ algebra (A, M) can be considered as an $\mathsf{E_{d-1}}$ algebra by forgetting the structure maps in equation 38. For the final adjunction, any $\mathsf{E_{d-1}}$ algebra in $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ has an underlying pair of $\mathcal{C}$-objects (A, M).

Notation 33. Throughout the remainder of this paper $\mathcal{C}$ will be a cofibrantly generated symmetric monoidal model category tensored over Top such that the adjunctions in definition 32 are Quillen adjunctions.

We use the model structure on the category of $\mathsf{E_{d-1}}$-A modules to define Hochschild cohomology.

Definition 34. Given an $\mathsf{E_{d-1}}$ algebra $A \in \mathcal{C}$, let the *Hochschild cohomology object* of A be

$$\mathrm{Hoch}(A) = \underline{\mathrm{hom}}^{A}_{\mathsf{E_{d-1}}}(A^c, A^f),$$

where $\underline{\mathrm{hom}}^{A}_{\mathsf{E_{d-1}}}$ is given by the equalizer in definition 30. The $\mathsf{E_{d-1}}$-A modules A^c and A^f are cofibrant and fibrant replacements for A respectively. Note that $\mathrm{Hoch}(A)$ is an object of $\mathcal{C}$.

We will use the degree 0 and 1 pieces of the swiss cheese operad to build a cofibrant replacement for A as an $\mathsf{E_{d-1}}$-A module.

3.2 Swiss cheese in degrees zero and one

An $\mathsf{SC_d^{h1}}$ algebra (see definition 32 and example 22) is a pair (A, M) of objects of $\mathcal{C}$ together with maps in $\mathrm{Coll}^{\leq 1}(\mathcal{C})$ for every m,

$$\big(\mathsf{E_{d-1}}(m), \mathsf{SC_d^h}(1, m)\big) \otimes (A, M)^{\otimes m} \to (A, M), \tag{35}$$

where we have used the isomorphism $\mathsf{SC_d^h}(0, m) \simeq \mathsf{E_{d-1}}(m)$. Alternatively, we can view the morphism in equation 35 as three separate maps in $\mathcal{C}$.

$$\mathsf{E_{d-1}}(m) \otimes A^{\otimes m} \to A \tag{36}$$
$$\mathsf{E_{d-1}}(m) \otimes M \otimes A^{\otimes m-1} \to M \tag{37}$$
$$\mathsf{SC_d^h}(1, m) \otimes A^{\otimes m} \to M \tag{38}$$

The condition that the maps in equation 35 define an $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ structure on the pair (A, M) is the condition that the diagram 39 commutes in the $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$-enriched category $\mathrm{Coll}^{\leq 1}(\mathcal{C})$.

$$\begin{tikzcd}
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}(m') \otimes \mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}(m; m') \otimes (A, M)^{\otimes m} \arrow[r] \arrow[d] & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}(m) \otimes (A, M)^{\otimes m} \arrow[d] \\
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}(m') \otimes (A, M)^{\otimes m'} \arrow[r] & (A, M)
\end{tikzcd} \tag{39}$$

In terms of equations 36, 37, and 38, diagram 39 splits into four diagrams. Each diagram is determined by the degrees of the three tensor factors in the upper left hand corner of diagram 39. In the first, diagram 40, the degrees are $0, 0, 0$. In 41, the degrees are $0, 0, 1$; in 42, degrees are $1, 0, 0$; and in 43, degrees are $0, 1, 0$.

$$\begin{tikzcd}
\mathsf{E}_{\mathsf{d-1}}(m') \otimes \mathsf{E}_{\mathsf{d-1}}(m; m') \otimes A^{\otimes m} \arrow[r] \arrow[d] & \mathsf{E}_{\mathsf{d-1}}(m) \otimes A^{\otimes m} \arrow[d] \\
\mathsf{E}_{\mathsf{d-1}}(m') \otimes A^{\otimes m'} \arrow[r] & A
\end{tikzcd} \tag{40}$$

The above diagram, 40, commutes for all m', m if and only if A is an $\mathsf{E}_{\mathsf{d-1}}$ algebra. Diagram 41 below commutes if and only if M is an $\mathsf{E}_{\mathsf{d-1}}$-$A$ module.

$$\begin{tikzcd}
\mathsf{E}_{\mathsf{d-1}}(m') \otimes \mathsf{E}_{\mathsf{d-1}}(m; m') \otimes M \otimes A^{\otimes m-1} \arrow[r] \arrow[d] & \mathsf{E}_{\mathsf{d-1}}(m) \otimes M \otimes A^{\otimes m'-1} \arrow[d] \\
\mathsf{E}_{\mathsf{d-1}}(m) \otimes M \otimes A^{\otimes m-1} \arrow[r] & M
\end{tikzcd} \tag{41}$$

Diagram 42 shows a compatibility condition between the degree 0 and degree 1 structures.

$$\begin{tikzcd}
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m') \otimes \mathsf{E}_{\mathsf{d-1}}(m; m') \otimes A^{\otimes m} \arrow[r] \arrow[d] & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m) \otimes A^{\otimes m} \arrow[d] \\
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m') \otimes A^{\otimes m'} \arrow[r] & M
\end{tikzcd} \tag{42}$$

Diagram 43 presents another compatibility condition between the degree 0 and degree 1 structures.

$$\begin{tikzcd}
\mathsf{E}_{\mathsf{d-1}}(m') \otimes \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m; 0, m') \otimes A^{\otimes m} \arrow[r] \arrow[d] & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m) \otimes A^{\otimes m} \arrow[d] \\
\mathsf{E}_{\mathsf{d-1}}(m') \otimes M \otimes A^{\otimes m'-1} \arrow[r] & M
\end{tikzcd} \tag{43}$$

Example 44. In this example we construct the universal extension of an $\mathsf{E}_{\mathsf{d-1}}$ algebra A to an $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ algebra. We denote this universal pair by (A, A^{sc}). The composite forgetful functor

$$\mathrm{Alg}_{\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}}(\mathrm{Coll}^{\leq 1}(\mathcal{C})) \to \mathrm{Alg}_{\mathsf{E}_{\mathsf{d-1}}}(\mathrm{Coll}^{\leq 1}(\mathcal{C})) \to \mathrm{Alg}_{\mathsf{E}_{\mathsf{d-1}}}(\mathcal{C}),$$

has a left adjoint which sends the $\mathsf{E}_{\mathsf{d-1}}$ algebra A to the pair (A, A^{sc}) where A^{sc}, which may be read as "A swiss cheese", is a quotient of

$$\bar{A}^{sc} = \coprod_{m \geq 0} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m) \otimes_{S_m} A^{\otimes m}, \tag{45}$$

where $S_m = \mathrm{Aut}(\underline{m})$. We can think of $\bar{A}^{sc}$ heuristically as $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, \bullet) \otimes A^{\otimes \bullet}$. Now both A and $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, \bullet)$ carry an action of $\mathsf{E}_{\mathsf{d-1}}$, so we can form the quotient $A^{sc} := \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, \bullet) \otimes_{\mathsf{E}_{\mathsf{d-1}}} A^{\otimes \bullet}$. More precisely, A^{sc} is defined as the coequalizer

$$\coprod_{m, m'} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m) \otimes \mathsf{E}_{\mathsf{d-1}}(m'; m) \otimes A^{\otimes m'} \rightrightarrows \bar{A}^{sc} \xrightarrow{\mathrm{coeq}} A^{sc}, \tag{46}$$

where one of the arrows is given by the operadic composition on swiss cheese and the other by the $\mathsf{E}_{\mathsf{d-1}}$ structure on A. Figure 2 shows the relation $\sim$ such that $A^{sc} = \bar{A}^{sc}/\sim$.



Figure 2: The relations in A^{sc} come from the $\mathsf{E}_{\mathsf{d-1}}$ algebra structure of A. If m is the map $A^{\otimes 3} \to A$ given by the swiss cheese element in $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(0,3) \simeq \mathsf{E}_{\mathsf{d-1}}(3)$ in the figure, set $a = m(a_1, a_2, a_3)$. The edges t_1 and t_2 are less than ∞, so the relation does not apply to the vertices on the left and right.

Verifying that (A, A^{sc}) is an $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ algebra is a matter of using the commuting left and right actions of $\mathsf{E}_{\mathsf{d-1}}$ on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$. By this we mean the morphism below uses both left and right actions, and can be obtained by performing the left action first, then the right, or vice versa,

$$\mathsf{E}_{\mathsf{d-1}}(m'') \times \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m'; 0, m'') \times \mathsf{E}_{\mathsf{d-1}}(m; m') \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(1, m).$$

The left action defines a map

$$\mathsf{E}_{\mathsf{d-1}}(m) \otimes \bar{A}^{sc} \otimes A^{\otimes m-1} \to \bar{A}^{sc}, \tag{47}$$

using the the $\circ_1$ operad composition. Since the left and right $\mathsf{E}_{\mathsf{d-1}}$ actions on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ commute, the arrow in 47 descends to give the data of a $\mathsf{E}_{\mathsf{d-1}}$-$A$ module structure on A^{sc}, i.e. equation 37 for $M = A^{sc}$. Of course, the maps from equation 38 with $M = A^{sc}$ are simply given by $\bar{A}^{sc} \to A^{sc}$.

Now let us observe that the four diagrams 40-43 commute for (A, A^{sc}). The first diagram, 40, is trivial since A is an $\mathsf{E}_{\mathsf{d-1}}$ algebra. The second diagram, 41, commutes since the left action of $\mathsf{E}_{\mathsf{d-1}}$ on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ is indeed an action. That is, it is compatible with composition in $\mathsf{E}_{\mathsf{d-1}}$. The third diagram, 42, clearly commutes. Indeed, this diagram is the reason the coequalizer $\bar{A}^{sc} \to A^{sc}$ in equation 46 was defined in the first place. Finally, the fourth diagram, 43, certainly commutes if $M = \bar{A}^{sc}$. In this case, note that the bottom map in diagram 43 corresponds to equation 47. Thus if we pass from $\bar{A}^{sc}$ to A^{sc} this last diagram still commutes since, by definition, the $\mathsf{E}_{\mathsf{d-1}}$-$A$ module structure on A^{sc} is defined using the quotient map $\bar{A}^{sc} \to A^{sc}$ together with equation 47.

Lemma 48. Fix an $\mathsf{E}_{\mathsf{d-1}}$ algebra A and consider A^{sc} as an $\mathsf{E}_{\mathsf{d-1}}$-$A$ module, then there is an isomorphism of categories

$$\mathrm{Mod}^{A}_{\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}}(\mathcal{C}) \cong \mathrm{Mod}^{A}_{\mathsf{E}_{\mathsf{d-1}}}(\mathcal{C})_{A^{sc}/},$$

where $\mathrm{Mod}^{A}_{\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}}(\mathcal{C})$ is the fiber over A of the forgetful functor $\mathrm{Alg}_{\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}}(\mathrm{Coll}^{\leq 1}(\mathcal{C})) \to \mathrm{Alg}_{\mathsf{E}_{\mathsf{d-1}}}(\mathcal{C})$

Proof. Let (A, M) be an $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h1}}$ algebra extending the existing $\mathsf{E}_{\mathsf{d-1}}$ algebra structure on A. Then $M \in \mathrm{Mod}^{A}_{\mathsf{E}_{\mathsf{d-1}}}(\mathcal{C})$ and the structure maps in equation 38, when combined for all m, give a map $\bar{A}^{sc} \to M$. Diagram 42 shows that this descends to $A^{sc} \to M$ and diagram 43 shows that this latter arrow is an $\mathsf{E}_{\mathsf{d-1}}$-$A$ module map.

On the other hand if M is an $\mathsf{E}_{\mathsf{d-1}}$-$A$ module, then M is already equipped with the data of equation 37 making diagram 41 commute. If $A^{sc} \to M$ is a morphism of $\mathsf{E}_{\mathsf{d-1}}$-$A$ modules, then M is equipped with the data of equation 38. The diagram 42 commutes because of relation defining A^{sc} in equation 46. The diagram 43 commutes since $A^{sc} \to M$ is a morphism of $\mathsf{E}_{\mathsf{d-1}}$-$A$ modules. ☐

Corollary 49. Let A be a cofibrant $\mathsf{E_{d-1}}$ algebra, then A^{sc} is a cofibrant $\mathsf{E_{d-1}}$-A module.

Proof. The forgetful functor $\mathrm{Alg}_{\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}}(\mathcal{C}) \to \mathrm{Alg}_{\mathsf{E_{d-1}}}(\mathcal{C})$ preserves fibrations (see definition 32). Thus the left adjoint of this forgetful functor, applied to the cofibrant object A, gives a cofibrant $\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}$ algebra (A, A^{sc}). Thus A^{sc} is cofibrant in $\mathrm{Mod}^{A}_{\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}}(\mathcal{C})$. Lemma 48 shows that the forgetful functor $\mathrm{Mod}^{A}_{\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}}(\mathcal{C}) \to \mathrm{Mod}^{A}_{\mathsf{E_{d-1}}}(\mathcal{C})$ preserves pushouts. The model structures here are cofibrantly generated, so A^{sc} is also cofibrant as an object of $\mathrm{Mod}^{A}_{\mathsf{E_{d-1}}}(\mathcal{C})$. ☐

Definition 50. Let $p(m) : \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \to \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(0, m) \simeq \mathsf{E_{d-1}}(m)$ be the projection which forgets the single full disc. We can make $p = (p(m))_{m\geq 0}$ into a morphism of operads in the following way. For each m consider the degree 0-1 collection $(\mathsf{E_{d-1}})^{\leq 1}(m) := (\mathsf{E_{d-1}}(m), \mathsf{E_{d-1}}(m))$. The structure of $\mathsf{E_{d-1}}$ as an operad in Top can be used to make $(\mathsf{E_{d-1}})^{\leq 1}$ an operad in $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$. This makes $(\mathsf{id}, p)\colon \mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}} \to (\mathsf{E_{d-1}})^{\leq 1}$ into a morphism of operads.

If A is an $\mathsf{E_{d-1}}$ algebra we can define a morphism in $\mathcal{C}$,

$$\bar{A}^{sc} = \coprod_{m\geq 0} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes_{S_m} A^{\otimes m} \to \coprod_{m\geq 0} \mathsf{E_{d-1}}(m) \otimes_{S_m} A^{\otimes m} \to A,$$

where the first arrow uses p and the second arrow uses the $\mathsf{E_{d-1}}$ algebra structure on A. This map factors to give a morphism of $\mathsf{E_{d-1}}$-A modules $p_A\colon A^{sc} \to A$.

Remark 51. By [Spi01, section 5] we can conclude that $\mathsf{Op}(\mathrm{Coll}^{\leq 1}(\mathsf{Top}))$ is tensored over Top. If $\mathcal{O} = (\mathcal{O}^0, \mathcal{O}^1)$ is an operad in degree 0-1 collections of topological spaces, and $K \in \mathsf{Top}$, then $K \otimes \mathcal{O}$ is defined to be the coequalizer

$$F(F(K \otimes \mathcal{O})) \rightrightarrows F(K \otimes \mathcal{O}) \xrightarrow{\text{coeq}} K \otimes \mathcal{O},$$

where $K \otimes \mathcal{O}$, is the collection of degree 0-1 collections whose arity m, degree i component is $K \times \mathcal{O}^i(m)$, and F is the free operad functor.

If $\mathcal{O}$ is a cofibrant operad in $\mathrm{Coll}^{\leq 1}(\mathsf{Top})$, then $[0, 1] \otimes \mathcal{O}$ is a cylinder object, and a homotopy $h\colon [0, 1] \otimes \mathcal{O} \to \mathcal{P}$ gives the data of maps $h^i(m)\colon [0, 1] \times \mathcal{O}^i(m) \to \mathcal{P}^i(m)$, which assemble into $h^i(m'; m)\colon [0, 1] \times \mathcal{O}^i(m'; m) \to \mathcal{P}(m'; m)$ for $i = 0, 1$ and $m', m \geq 0$. The homotopy h is compatible with operad composition in the sense that if $\alpha \in \mathcal{O}^i(m)$ and $\beta \in \mathcal{O}^j(m'; m)$, $i + j \leq 1$, and $t \in [0, 1]$, then

$$h^i(m)(t, \alpha) \circ h^j(m'; m)(t, \beta) = h^{i+j}(m')(t, \alpha \circ \beta).$$

Lemma 52. For any $\mathsf{E_{d-1}}$ algebra $A \in \mathcal{C}$, the map $A^{sc} \to A$ is a weak equivalence of $\mathsf{E_{d-1}}$-A modules.

Proof. Abusing notation we write $p := (\mathsf{id}, p)$ for the morphism of operads from definition 50. One can show that p is a weak equivalence of fibrant and cofibrant operads in degree 0-1 collections of topological spaces. Therefore there is a map of operads $\iota\colon (\mathsf{E_{d-1}})^{\leq 1} \to \mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}$ and there are homotopies $h\colon \mathsf{id}_{\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}} \simeq \iota p$, and $g\colon \mathsf{id}_{(\mathsf{E_{d-1}})^{\leq 1}} \simeq p\iota$. Since $[0, 1] \otimes -$ distributes over coequalizers, we can use h to define a homotopy $h_A\colon [0, 1] \otimes A^{sc} \to A^{sc}$,

$$\begin{tikzcd}
\coprod_m [0,1] \otimes \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes A^{\otimes m} \arrow[r] \arrow[d, "\sqcup h^1(m) \otimes \mathsf{id}_A^{\otimes m}"] & {[0,1] \otimes A^{sc}} \arrow[d, "h_A"] \\
\coprod_m \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes A^{\otimes m} \arrow[r] & A^{sc},
\end{tikzcd}$$

where $h^1(m)$ is defined from h as in remark 51. When $t = 1$, $h_A(1, -)$ factors as

$$\begin{tikzcd}
\coprod_m \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1,m) \otimes A^{\otimes m} \arrow[r] \arrow[d] & A^{sc} \arrow[d, "p_A"] \\
\coprod_m \mathsf{E}_{\mathsf{d-1}}(m) \otimes A^{\otimes m} \arrow[r] \arrow[d] & A \arrow[l, dashed, bend right, "s"'] \arrow[d, "\iota_A"] \\
\coprod_m \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1,m) \otimes A^{\otimes m} \arrow[r] & A^{sc},
\end{tikzcd}$$

where p_A is the map from definition 50, and ι_A is the evident composite in the diagram using s. The map s is a section of the middle horizontal arrow, defined using the identity of the operad $\mathsf{E}_{\mathsf{d-1}}$,

$$A \simeq \{1_{\mathsf{E}_{\mathsf{d-1}}}\} \otimes A \to \mathsf{E}_{\mathsf{d-1}}(1) \otimes A \to \coprod_m \mathsf{E}_{\mathsf{d-1}}(m) \otimes A^{\otimes m}.$$

The map ι_A is a map of $\mathsf{E}_{\mathsf{d-1}}$-$A$ modules, and h_A is a homotopy $\mathsf{id}_{A^{sc}} \simeq \iota_A p_A$. Similarly, g defines a homotopy $g_A \colon \mathsf{id}_A \to p_A \iota_A$. $\square$

The precise sense in which Hochschild cohomology can be obtained from the degree 0-1 pieces of the swiss cheese operad is contained in the following

Proposition 53. Let A be a fibrant and cofibrant $\mathsf{E}_{\mathsf{d-1}}$ algebra. Then the Hochschild cohomology object of A can be computed as

$$\mathrm{Hoch}(A) \simeq \underline{\mathrm{hom}}^A_{\mathsf{E}_{\mathsf{d-1}}}(A^{sc}, A).$$

Proof. We are using the projective model structure (definition 32), so A is fibrant as an object of $\mathcal{C}$ and thus as a $\mathsf{E}_{\mathsf{d-1}}$-$A$ module. By corollary 49 and lemma 52, A^{sc} is a cofibrant replacement for A as an $\mathsf{E}_{\mathsf{d-1}}$-$A$ module. By definition 34, this proves the proposition. $\square$

4 The universal cheese theorem

In the one-colored operad $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ the single full disc was never considered as input, only as a marker of degree one. Allowing the single disc to be considered as giving an input means viewing $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ as a partially defined 2-colored operad. Rather than making the notion *partially defined* precise, we simply define the notion of a 2-colored algebra over $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ in definition 54. Proposition 55 is a version of the swiss cheese theorem for the operad $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$. That is, a 2-colored $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ structure on the pair (B, A) is equivalent to a $\mathcal{C}$-morphism, $B \to \mathrm{Hoch}(A)$. In other words, $\mathrm{Hoch}(A)$ is the universal object of $\mathcal{C}$ acting on the $\mathsf{E}_{\mathsf{d-1}}$ algebra A through $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$. This result is generalized twice, first in corollary 62, then in proposition 75. We refer to proposition 75 as the universal swiss cheese theorem since it replaces E_d in the swiss cheese theorem with an arbitrary operad.

Definition 54. Let A be an $\mathsf{E}_{\mathsf{d-1}}$ algebra. Let $\mathcal{C}_{/\mathrm{Hoch}(A)}$ denote the over category of $\mathrm{Hoch}(A) \in \mathcal{C}$. More precisely, the objects are $\mathcal{C}$-morphisms $B \to \mathrm{Hoch}(A)$ and the morphisms are $\mathcal{C}$ morphisms $B \to B'$ commuting with the maps to $\mathrm{Hoch}(A)$. In addition, let $\mathrm{Alg}^A_{\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}}(\mathcal{C})$ denote the category of $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ algebras of the form (B, A) where the induced $\mathsf{E}_{\mathsf{d-1}}$ structure on A is the one given. Morphisms are maps of $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$ algebras which are identity on A.

Given a pair of objects (B, A) of $\mathcal{C}$, we let $\mathrm{End}^{\mathrm{h1}}(B, A)$ denote the operad obtained by applying the forgetful functor $\mathsf{Op}_K \to \mathsf{Op}(\mathrm{Coll}^{\leq 1})$ to $\mathrm{End}(B, A)$ from 11 and 12.

Let $\mathrm{Alg}^A_{\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}}(\mathcal{C})$ denote the category of objects $B \in \mathcal{C}$ together with a morphism in $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$, $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}} \to \mathrm{End}^{\mathrm{h1}}(B, A)$ extending the $\mathsf{E}_{\mathsf{d-1}}$ structure on A. A morphism $B \to B'$ in $\mathcal{C}$ induces a map of operads $\mathrm{End}^{\mathrm{h1}}(B', A) \to \mathrm{End}^{\mathrm{h1}}(B, A)$. Such a morphism gives a map in $\mathrm{Alg}^A_{\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}}(\mathcal{C})$ if this induced map respects to maps from $\mathsf{SC}^{\mathrm{h1}}_{\mathsf{d}}$.

Proposition 55. There is an isomorphism of categories

$$\mathrm{Alg}^{A}_{\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}}(\mathcal{C}) \cong \mathcal{C}_{/\mathrm{Hoch}(A)}.$$

Proof. The data of an algebra on the left hand side is an object $B \in \mathcal{C}$ together with maps

$$\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes B \otimes A^{\otimes m} \to A, \tag{56}$$

for each $m \geq 0$. The conditions on 56 are that diagrams 57 and 58 commute.

$$\begin{tikzcd}
\mathsf{E}_{\mathsf{d-1}}(m') \otimes \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m; 0, m') \otimes B \otimes A^{\otimes m} \arrow[r] \arrow[d] & \mathsf{E}_{\mathsf{d-1}}(m') \otimes A^{\otimes m'} \arrow[d] \\
\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes B \otimes A^{\otimes m} \arrow[r] & A,
\end{tikzcd} \tag{57}$$

$$\begin{tikzcd}
\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m') \otimes \mathsf{E}_{\mathsf{d-1}}(m; m') \otimes B \otimes A^{\otimes m} \arrow[r] \arrow[d] & \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m) \otimes B \otimes A^{\otimes m} \arrow[d] \\
\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m') \otimes B \otimes A^{\otimes m'} \arrow[r] & A.
\end{tikzcd} \tag{58}$$

Equivalently, we can use the hom-tensor adjunction and assemble the maps in 56 to a single map $B \to \underline{\mathrm{hom}}_{\mathcal{C}}(\bar{A}^{sc}, A)$ (see 45). The commutativity of diagram 58 is equivalent to this map lifting to

$$B \to \underline{\mathrm{hom}}_{\mathcal{C}}(A^{sc}, A). \tag{59}$$

Note that, dual to 46, $\underline{\mathrm{hom}}_{\mathcal{C}}(A^{sc}, A)$ is given by the equalizer

$$\underline{\mathrm{hom}}_{\mathcal{C}}(A^{sc}, A) \xrightarrow{\mathrm{eq}} \underline{\mathrm{hom}}_{\mathcal{C}}(\bar{A}^{sc}, A) \rightrightarrows \underline{\mathrm{hom}}_{\mathcal{C}}(\coprod_{m,m'} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(1, m') \otimes \mathsf{E}_{\mathsf{d-1}}(m; m') \otimes A^{\otimes m}, A).$$

With this observation we can now rewrite diagram 57 as

$$\begin{tikzcd}
B \otimes F^{A}_{\mathsf{E}_{\mathsf{d-1}}}(A^{sc}) \arrow[r] \arrow[d] & F^{A}_{\mathsf{E}_{\mathsf{d-1}}}(A) \arrow[d] \\
B \otimes A^{sc} \arrow[r] & A.
\end{tikzcd} \tag{60}$$

Recall that $F^{A}_{\mathsf{E}_{\mathsf{d-1}}} \colon \mathcal{C} \to \mathcal{C}$ is the free $\mathsf{E}_{\mathsf{d-1}}$-$A$ module functor, see definition 30. Clearly, diagram 57 commutes if and only if diagram 60 commutes and if and only if the map 59 factors through $\mathrm{Hoch}(A) = \underline{\mathrm{hom}}^{A}_{\mathsf{E}_{\mathsf{d-1}}}(A^{sc}, A)$. We conclude that the data of an $\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}$ algebra structure on (B, A) is the data of an $\mathsf{E}_{\mathsf{d-1}}$ algebra structure on A together with a $\mathcal{C}$-morphism $B \to \mathrm{Hoch}(A)$.

It is clear that a map $(B, A) \to (B', A)$ which is identity on A gives an $\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}$ algebra morphism if and only if the map $B \to B'$ commutes with the corresponding morphisms to $\mathrm{Hoch}(A)$.

□

Definition 61. Define $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}} \in \mathsf{Op}(\mathrm{Coll})$ as the image of the left adjoint of $\mathsf{Op}(\mathrm{Coll}) \to \mathsf{Op}(\mathrm{Coll}^{\leq 1})$ applied to $\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}}$. Following definition 54, define $\mathrm{Alg}^{A}_{\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}}(\mathcal{C})$ as the category of objects $B \in \mathcal{C}$ together with a morphism in $\mathsf{Op}(\mathrm{Coll})$, $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}} \to \mathrm{End}^{\mathrm{h}}(B, A)$ extending the $\mathsf{E}_{\mathsf{d-1}}$ structure on A.

Corollary 62. There is an isomorphism of categories

$$\mathrm{Alg}^{A}_{\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}}(\mathcal{C}) \cong \mathcal{C}_{/\mathrm{Hoch}(A)}.$$

Proof. The adjunction isomorphism puts operad maps $\mathsf{SC}^{\mathrm{h}1}_{\mathsf{d}} \to \mathrm{End}^{\mathrm{h}1}(B, A)$ in one-to-one correspondence with operad maps $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}} \to \mathrm{End}^{\mathrm{h}}(B, A)$. □

Definition 63. For each $n \geq 0$, let $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, \bullet)$ denote the operad in $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$ whose arity m component is the degree 0-1 collection $(\mathsf{E}_{\mathsf{d-1}}(m), \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, m))$. One may think of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, \bullet)$ as a bimodule over $\mathsf{E}_{\mathsf{d-1}}$. Let $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ be the collection whose n^{th} space is $\mathsf{E}_{\mathsf{d-1}}$ bimodule maps

$$\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(n) \coloneqq \mathrm{map}_{\mathsf{E}_{\mathsf{d-1}}}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1, \bullet), \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, \bullet)).$$

Lemma 64. Operadic composition in $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ induces the structure of an operad on the collection $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$.

Proof. Define an operad $\mathcal{E} \in \mathsf{Op}(\mathrm{Coll})$ by setting

$$\mathcal{E}(n, m) = \mathrm{map}_{\mathrm{Coll}}\big(\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(\bullet; n),\ \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, m)\big).$$

Let $n_0 + \cdots + n_m = n$ and $\ell_1 + \cdots + \ell_m = \ell$. The monoidal structure $\otimes$ on collections gives a map from $\mathcal{E}(n_0, m) \times \prod_{i=1}^{m} \mathcal{E}(n_i, \ell_i)$ to

$$\mathrm{map}_{\mathrm{Coll}}\left(\bigotimes_{i=0}^{m} \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(\bullet; n_i),\ \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, m) \otimes \Big(\bigotimes_{i=1}^{m} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, \ell_i)\Big)\right). \tag{65}$$

Now push forward from 65 via the operad structure on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, m) \otimes \Big(\bigotimes_{i=1}^{m} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, \ell_i)\Big) \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(\bullet, \ell),$$

and pull back from 65 by

$$\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(\bullet; n) \to \bigotimes_{i=0}^{m} \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(\bullet; n_i). \tag{66}$$

This defines

$$\mathcal{E}(n_0, m) \times \prod_{i=1}^{m} \mathcal{E}(n_i, \ell_i) \to \mathcal{E}(n, \ell).$$

The morphism in 66 comes from the sequence of maps

$$\begin{aligned}
\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(k; n) &= \coprod_{f\colon \underline{k} \to \underline{n}} \prod_{i=1}^{n} \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(f^{-1}(i)) && (67)\\
&= \coprod_{f\colon \underline{k} \to \underline{n}} \prod_{j \in \underline{0} \sqcup \underline{m}} \prod_{i \in g^{-1}(j)} \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(f^{-1}(i)) && (68)\\
&\to \coprod_{\tilde{f}\colon \underline{k} \to \underline{0} \sqcup \underline{m}} \prod_{j \in \underline{0} \sqcup \underline{m}} \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(\tilde{f}^{-1}(j); n_j). && (69)
\end{aligned}$$

The first equality holds by definition, the second is a regrouping. The decomposition $\sum_{i=0}^{m} n_i = n$ defines a map $g\colon \underline{n} \to \underline{0} \sqcup \underline{m}$ where $\left|g^{-1}(i)\right| = n_i$. The third map sends the component corresponding to $f\colon \underline{k} \to \underline{n}$ to the component corresponding to $fg\colon \underline{k} \to \underline{0} \sqcup \underline{m}$. For each n, m there is a map,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, m) \to \mathcal{E}(n, m). \tag{70}$$

When $n = 0, 1$, the map 70 is canonical. Restricting to degrees 0 and 1 gives a map of $\mathsf{Op}(\mathrm{Coll}^{\leq 1})$ operads $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}1} \to \mathcal{E}^1$, where $\mathcal{E}^1$ is the degree 0-1 part of $\mathcal{E}$. Since $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ is freely generated by its degree 0 and 1 pieces, we get 70 for all n, assembling into a map of operads in $\mathsf{Op}(\mathrm{Coll})$. This guarantees that 71 can be used to define an operadic composition law on $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n, m) \times \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(k_1) \times \cdots \times \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(k_n) \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(k, m). \tag{71}$$

□

Definition 72. Let $\mathcal{O}$ be any 1-colored topological operad, and let $\rho\colon \mathcal{O} \to \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ be a map of operads. Define the K-colored operad $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O}$ by setting

$$(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O})^{\mathrm{h}}(n,m) = \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m) \qquad (\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O})^{\mathrm{f}}(n,m) = \begin{cases} \mathcal{O}(n) & m = 0 \\ \emptyset & m > 0. \end{cases}$$

Composition in $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O}$ uses composition in $\mathcal{O}$, composition in $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$, and the action of $\mathcal{O}$ on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ defined by ρ,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m) \times \mathcal{O}(k;n) \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m) \times \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(k;n) \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(k,m),$$

where the right arrow above is the one in 71.

Lemma 73. *Let $\mathcal{O}$ be a topological operad and let $\rho\colon \mathcal{O} \to \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ be a map of operads, then the $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ structure on (H,A) naturally extends to a $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O}$ structure on (H,A). In particular, $H = \mathrm{Hoch}(A)$ inherits an $\mathcal{O}$ algebra structure.*

Proof. We only need to show there is a map of operads $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}) \to \mathrm{End}(H)$ compatible with the action of $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ and the action of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ on H. Indeed, the map

$$\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(n) \otimes H^{\otimes n} \to H \tag{74}$$

is adjoint to the maps, for all $m \geq 0$,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1,m) \otimes \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(n) \otimes H^{\otimes n} \otimes A^{\otimes m} \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m) \otimes H^{\otimes n} \otimes A^{\otimes m} \to A,$$

where the first arrow is 71 and the second arrow is the $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ structure on (H,A). To check that 74 is compatible with composition in $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ observe that there are two morphisms of operads in $\mathsf{Op}(\mathrm{Coll})$,

$$\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rightrightarrows \mathrm{End}^{\mathrm{h}}(\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(H), A),$$

where $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(H)$ is the free $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ algebra generated by H. One of the arrows uses the action of $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$, while the other uses the map $\mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})(H) \to H$ defined by 74. To check that these arrows agree, we only need to check that they agree out of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m)$ when $n = 0,1$. This is because $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ is freely generated in degrees 0 and 1. When $n = 0$, the maps are obviously the same. When $n = 1$, the maps are the same by definition of the $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}1}$ structure on (H,A). $\square$

Proposition 75 (The universal cheese conjecture)**.** *Let $\mathcal{O}$ be a topological operad and let $\rho\colon \mathcal{O} \to \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$ be a map of operads. Then using the induced $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O}$ structure on $(\mathrm{Hoch}(A), A)$ from lemma 73 gives an isomorphism of categories*

$$\mathrm{Alg}^{A}_{(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O})}(\mathcal{C}) \cong \mathrm{Alg}_{\mathcal{O}}(\mathcal{C})_{/\mathrm{Hoch}(A)}.$$

Proof. Given any $\mathcal{C}$ morphism $B \to H$ we can form the following diagram. For brevity, we have deleted appearances $\otimes$.

$$\begin{tikzcd}
 & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1,m)\mathcal{O}(n)B^nA^m \arrow[rr] \arrow[dl] \arrow[dd] & & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m)B^nA^m \arrow[dl] \arrow[dd] \\
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1,m)\mathcal{O}(n)H^nA^m \arrow[rr, crossing over] \arrow[dd] & & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(n,m)H^nA^m & \\
 & \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1,m)BA^m \arrow[rr] \arrow[dl] & & A \arrow[dl, "="] \\
\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}(1,m)HA^m \arrow[rr] & & A \arrow[from=uu, crossing over] &
\end{tikzcd} \tag{76}$$

Let (B,A) be a $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes_\rho \mathcal{O}$ algebra extending the given $\mathsf{E}_{\mathsf{d-1}}$ structure on A, then by corollary 62 we get a $\mathcal{C}$ morphism $B \to H = \mathrm{Hoch}(A)$ making the right face of the cube 76 commute. The front face commutes by lemma 73. The back face commutes by assumption. The bottom face commutes by definition, and the top face commutes trivially. This implies that, after composition with the maps whose codomain is A, the left face of the cube commutes. By adjointness, the two maps $\mathcal{O}(B) \rightrightarrows H$ agree, implying that $B \to H$ is indeed an $\mathcal{O}$ algebra morphism.

On the other hand, given an $\mathcal{O}$ algebra B together with an $\mathcal{O}$ algebra map $B \to H$, we get an $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$ structure on (B, A) from the underlying $\mathcal{C}$ morphism. We only need to check that the $\mathcal{O}$ structure on B and the $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$ structure on (B, A) are compatible via ρ. Indeed, since $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$ is freely generated in degrees 0 and 1, it is enough to check that the back face of the cube commutes. But this holds because all other faces commute. Most importantly, the left face commutes because $B \to H$ is an $\mathcal{O}$ algebra map.

It is easy to see that each of these constructions are natural in B and are inverse to one another. □

5 The homotopy E_d structure on $\mathrm{Hoch}(A)$

In light of proposition 75, to prove the swiss cheese theorem (15) we need to construct $\mathsf{E} \simeq E_d$ and an operad morphism $\mathsf{E} \to \mathrm{End}(\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty})$ in such a way that the corresponding K-colored operad $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \rtimes \mathsf{E}$ is equivalent to SC_d. While $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$ has no obvious action of E_d, it is equivalent to something that does have an E_d action. The following is in [Tho12, section 3].

Theorem 77. The natural map $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \to \mathsf{SC}_\mathsf{d}^{\mathrm{h}}$ is an acyclic cofibration of operads from $\mathsf{Op}(\mathrm{Coll})$.

In this section, we define the precise sense in which theorem 77 gives us our E_d action on $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$ up to homotopy. First, we have a lift p in the following diagram,

$$\begin{tikzcd}
\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \arrow[r, "\mathsf{id}"] \arrow[d, "\iota"'] & \mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty} \\
\mathsf{SC}_\mathsf{d}^{\mathrm{h}} \arrow[ur, dashed, "p"'] &
\end{tikzcd}$$

We know that $\mathsf{SC}_\mathsf{d}^{\mathrm{h}}$ is cofibrant since it is obtained as the W construction applied to a Σ-cofibrant, well-pointed operad SC_d^{h} [BM06]. Thus, by the corner axiom (Quillen's SM7) for monoidal model categories tensored over topological spaces [Spi01] we have an acyclic fibration

$$\mathrm{map}(\mathsf{SC}_\mathsf{d}^{\mathrm{h}}, \mathsf{SC}_\mathsf{d}^{\mathrm{h}}) \xrightarrow{\iota^*} \mathrm{map}(\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}, \mathsf{SC}_\mathsf{d}^{\mathrm{h}})$$

given by pre-composing with ι. Since both ιp and id live over ι, they must be homotopic. Let $h \colon [0, \infty] \otimes \mathsf{SC}_\mathsf{d}^{\mathrm{h}} \to \mathsf{SC}_\mathsf{d}^{\mathrm{h}}$ be a homotopy with $h_0 = \mathsf{id}$ and $h_\infty = \iota p$.

We will use this h to define a homotopy right E_d module structure on $\mathsf{SC}_\mathsf{d}^{\mathrm{h}\infty}$. For this we will use a homotopy equivalent version of E_d which sits inside the W construction. For simplicity we denote it by E. First we define the category LE_d. This category is not monoidal, but will be used to build E. The letter L stands for *level trees*. The objects of the topological category LE_d are finite sets, and the morphism space $LE_d(n, n')$ is defined to be a quotient of

$$\coprod_{\substack{k \geq 0 \\ n_1, \ldots, n_k}} E_d(n_1, n') \times E_d(n_2, n_1) \times \cdots \times E_d(n, n_k) \times [0, \infty]^k.$$

A point of the space above is given by a sequence $\alpha_i \in E_d(n_i, n_{i-1})$ for $1 \leq i \leq k+1$ and $t_i \in [0, \infty]$ for $1 \leq i \leq k$. For convenience of notation, we set $n_0 = n'$, $n_{k+1} = n$, $t_0 = \infty$, and $t_{k+1} = \infty$. We impose the following relations

Relations 78. If $t_i = 0$, then we can delete t_i and replace $(\ldots, \alpha_i, \alpha_{i+1}, \ldots)$ by the composition $(\ldots, \alpha_i \circ \alpha_{i+1}, \ldots)$. If $n_i = n_{i-1}$ and α_i is the identity, and $t_{i-1} = \infty = t_i$, then we can delete α_i from the sequence and delete t_i from the sequence. are defining t_0 and t_{k+1} to be ∞.

Remark 79. In the W construction, we could always delete the appearance of an identity and sum the lengths of the surrounding edges. We do not allow that here since we do not have $h_{s+t} = h_s \circ h_t$.

Composition in the category LE_d is given by concatenating sequences, setting the new coordinate in the factor $[0, \infty]$ between the two sequences to be ∞.

We can use the action of E_d on $\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}$ as well as the maps h_t, p, ι to define

$$LE_d(n, n') \to \mathrm{map}(\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n', m), \mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n, m)). \tag{80}$$

To do this, represent $\alpha \in LE_d(n, n')$ with a sequence $n' = n_0, n_1, \ldots, n_k, n_{k+1} = n$ together with $\alpha_i \in E_d(n_i, n_{i-1})$ for $1 \leq i \leq k+1$ and $t_i \in [0, \infty]$ for $1 \leq i \leq k$. This gives a chain of maps

$$\begin{aligned}\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n', m) \xrightarrow{\iota} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_0, m) \xrightarrow{\alpha_1} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_1, m) \xrightarrow{h_{t_1}} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_1, m) \xrightarrow{\alpha_2} \cdots \\ \cdots \xrightarrow{h_{t_k}} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_k, m) \xrightarrow{\alpha_{k+1}} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_{k+1}, m) \xrightarrow{p} \mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n, m).\end{aligned} \tag{81}$$

The maps $\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \xrightarrow{\alpha_i} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_{i+1}, m)$ are defined by the action of E_d on $\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}$,

$$\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \times E_d(n_{i+1}, n_i) \to \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_{i+1}, m).$$

Let us check that the relations 78 in LE_d are satisfied and that composition in E_d corresponds to composition of maps of $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}$. Suppose $t_i = 0$ for some i. Then $h_0 = \mathsf{id}$ so our chain of arrows contains

$$\mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_{i-1}, m) \xrightarrow{\alpha_i} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \xrightarrow{\alpha_{i+1}} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_{i+1}, m).$$

The composition of these two is equal to the map given by $\alpha_i \alpha_{i+1} \in E_d(\underline{n}_{i+1}, \underline{n}_{i-1})$. This is because $\mathsf{SC}^{\mathsf{h}}_{\mathsf{d}}(-, m)$ is a right E_d module.

If $n_i = n_{i-1}$, α_i is identity, and $t_{i-1} = t_i = \infty$, then the composition $h_{t_{i-1}} \circ \alpha_i \circ h_{t_i}$ is equal to $h_{t_{i-1}} = h_\infty$, so we are justified in deleting α_i and t_i from the sequence.

Now suppose we have some $t_i = \infty$, so that $\alpha \in LE_d(n, n')$ decomposes as $\beta_1 \beta_2$ for some $\beta_1 \in \mathsf{E}(n_i, n')$ and $\beta_2 \in \mathsf{E}(n, n_i)$. The chain of compositions defining the action of α from $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(1, m)$ to $\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n, m)$ contains the following segment.

$$\cdots \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \xrightarrow{h_\infty} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \to \cdots$$

The composite of the actions of β_1 and β_2 is computed by joining the chains for β_1 and for β_2. This joined chain agrees with the chain for α except for the segment above, which is replaced with the segment

$$\cdots \to \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \xrightarrow{p} \mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}(n_i, m) \xrightarrow{\iota} \mathsf{SC}^{\mathrm{h}}_{\mathsf{d}}(n_i, m) \to \cdots$$

Since $h_\infty = \iota p$, these chains of maps have the same composition.

The maps 80 define a functor

$$LE_d \to \mathrm{End}(\mathsf{SC}^{\mathrm{h}\infty}_{\mathsf{d}}). \tag{82}$$

There is no obvious operad structure on LE_d so we take the smallest operad containing LE_d. More precisely, 82 is a morphism of topological categories whose objects are finite sets. There is a forgetful functor from operads to the category of such topological categories. The operad E is defined to be the result of applying the left adjoint of this forgetful functor to the category LE_d.

Definition 83. Let $F(LE_d)$ be the free one-colored operad generated by the collection $n \mapsto LE_d(n,1)$. For each $n, n' \geq 0$ let $\mathsf{E}(n,n')$ be the topological space given by the coequalizer

$$\coprod_{n''} LE_d(n'',n') \times LE_d(n,n'') \rightrightarrows F(LE_d)(n,n') \xrightarrow{\mathrm{eq}} \mathsf{E}(n,n'),$$

where the two maps are given by composition in either $F(LE_d)$ or LE_d and the inclusion of LE_d into $F(LE_d)$.

Lemma 84. The category E is an operad and is equivalent to E_d.

Proof. Given a tree with its internal edges labeled by lengths $[0,\infty]$, call it a *level tree* if edges equidistant from the root vertex have the same length. Every morphism in $LE_d(n,1)$ can be represented by a level tree with vertices labeled by E_d. We can represent a point of $F(LE_d)$ with a tree whose vertices are labeled by level trees in LE_d. The relation defining $F(LE_d) \to \mathsf{E}$ allows us to break up a level tree with at least one level of length ∞ into several level trees all of whose levels have finite length. We conclude that E consists of trees labeled by E_d on the vertices, and $[0,\infty]$ on the internal edges, satisfying the condition that every maximal finite subtree is level.

There is an operad morphism $\mathsf{E} \to E_d$ which collapses all edge lengths to 0. On the level of collections, there is a homotopy inverse $E_d \to \mathsf{E}$. The homotopy $g_t \colon \mathsf{E} \to \mathsf{E}$ first collapses lengths of the edges furthest from the root to zero. This preserves the condition that every maximal finite subtree is level. Continuing in this way, we collapse all edge lengths to zero. □

The adjoint to 82 is an operad morphism $\mathsf{E} \to \mathrm{End}(\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty})$, which by definition 72 we can use to define the K-colored operad $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes \mathsf{E}$.

Lemma 85. The $\{\mathrm{f},\mathrm{h}\}$-colored operad $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes \mathsf{E}$ is weakly equivalent to the swiss cheese operad.

Proof. First, note that SC_d is equivalent to the semi-direct product of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ and E_{d} where the action of E_{d} factors through the map $\mathsf{E}_{\mathsf{d}} \to E_d$ which sends all lengths of internal edges to zero. This is because the map $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}} \to SC_d^{\mathrm{h}}$ which collapses trees is a weak equivalence and respects the action of E_{d}.

The action of E on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty}$ can be extended to an action on all of $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$. The sequence $\alpha_1, t_1, \cdots, t_k, \alpha_{k+1}$ acts via the composition

$$\begin{aligned}\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_0,m) \xrightarrow{\alpha_1} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_1,m) \xrightarrow{h_{t_1}} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_1,m) \to \cdots \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_{l-1},m) \\ \xrightarrow{h_{t_l}} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_{l-1},m) \xrightarrow{\alpha_l} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_l,m) \xrightarrow{h_\infty} \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}(n_l,m).\end{aligned} \tag{86}$$

Define for each $s \in [0,\infty]$ a homotopy $h^{[0,s]} \colon [0,\infty] \otimes \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}} \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ by setting $h_t^{[0,s]} = h_{\min(s,t)}$. We have $h_0^{[0,s]} = \mathsf{id}$ and $\mathrm{h}_\infty^{[0,s]} = h_s$, therefore we can define an action of E on $[0,\infty]_s \otimes \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ by replacing h_{t_i} in (86) with $h_{t_i}^{[0,s]}$. Then, when $s = 0$ each $h_{t_i}^{[0,0]}$ is the identity, so the action factors through the map $\mathsf{E} \to E_d$ collapsing all edges to 0. When $s = \infty$ we have $h_{t_i}^{[0,\infty]} = h_{t_i}$ so the action of E on $\mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}$ is 86. Thus we have a diagram of equivalences

$$SC_d \leftarrow \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}} \rtimes_{s=0} \mathsf{E} \to ([0,1] \otimes \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}}) \rtimes \mathsf{E} \leftarrow \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}} \rtimes_{s=\infty} \mathsf{E} \to \mathsf{SC}_{\mathsf{d}}^{\mathrm{h}\infty} \rtimes \mathsf{E},$$

where the map on the left collapses all edge lengths to 0. □

This proves the main theorem of the paper, 15.

References

[BF04] C. Berger and B. Fresse, *Combinatorial operad actions on cochains*, Math. Proc. Cambridge Philos. Soc. **137**(1), 135–174 (2004).

[BM03] C. Berger and I. Moerdijk, *Axiomatic homotopy theory for operads*, Comment. Math. Helv. **78**(4), 805–831 (2003), math/0206094v3.

[BM06] C. Berger and I. Moerdijk, *The Boardman-Vogt resolution of operads in monoidal model categories*, Topology **45**(5), 807–849 (2006), math/0502155v2.

[BV73] J. M. Boardman and R. M. Vogt, *Homotopy invariant algebraic structures on topological spaces*, Lecture Notes in Mathematics, Vol. 347, Springer-Verlag, Berlin, 1973.

[Coh76] F. R. Cohen, The homology of C_{n+1}-spaces, $n \geq 0$, in *The homology of iterated loop spaces*, Lecture Notes in Mathematics, no. 533, pages 207–351, Berlin-New York, 1976, Springer-Verlag.

[DTT09] V. Dolgushev, D. Tamarkin and B. Tsygan, *Proof of Swiss Cheese Version of Deligne's Conjecture*, ArXiv e-prints (April 2009), 0904.2753.

[Ger63] M. Gerstenhaber, *The cohomology structure of an associative ring*, Ann. of Math. (2) **78**, 267–288 (1963).

[Har07] J. E. Harper, *Homotopy theory of modules over operads and non-Sigma operads in monoidal model categories*, (2007), 0801.0191.

[HKV06] P. Hu, I. Kriz and A. A. Voronov, *On Kontsevich's Hochschild cohomology conjecture*, Compos. Math. **142**(1), 143–168 (2006), math/0309369v2.

[Hov99] M. Hovey, *Model categories*, volume 63 of *Mathematical Surveys and Monographs*, American Mathematical Society, Providence, RI, 1999.

[Kon99] M. Kontsevich, *Operads and motives in deformation quantization*, Lett. Math. Phys. **48**(1), 35–72 (1999), math/9904055, Moshé Flato (1937–1998).

[KS00] M. Kontsevich and Y. Soibelman, Deformations of algebras over operads and the Deligne conjecture, in *Conférence Moshé Flato 1999, Vol. I (Dijon)*, volume 21 of *Math. Phys. Stud.*, pages 255–307, Kluwer Acad. Publ., Dordrecht, 2000.

[Lur11] J. Lurie, *Higher algebra*, Prepublication book draft. Available from http://www.math.harvard.edu/lurie (2011).

[May72] J. May, *The geometry of iterated loop spaces*, Springer-Verlag, 1972.

[MS02] J. E. McClure and J. H. Smith, A solution of Deligne's Hochschild cohomology conjecture, in *Recent progress in homotopy theory (Baltimore, MD, 2000)*, volume 293 of *Contemp. Math.*, pages 153–193, Amer. Math. Soc., Providence, RI, 2002.

[MSS07] M. Markl, S. Shnider and J. Stasheff, *Operads in algebra, topology and physics*, volume 96, Amer Mathematical Society, 2007.

[Sin06] D. Sinha, *The homology of the little disks operad*, Arxiv preprint math/0610236 (2006).

[Spi01] M. Spitzweck, *Operads, Algebras and Modules in General Model Categories*, (2001), math/0101102.

[Tam98] D. E. Tamarkin, *Another proof of M. Kontsevich formality theorem*, (1998), math/9803025v4.

[Tho12] J. D. Thomas, *Forests and the W Construction*, (2012), math/1210.0883.

[Vor99] A. A. Voronov, The Swiss-cheese operad, in *Homotopy invariant algebraic structures (Baltimore, MD, 1998)*, volume 239 of *Contemp. Math.*, pages 365–373, Amer. Math. Soc., Providence, RI, 1999.

[Vor00] A. A. Voronov, Homotopy Gerstenhaber algebras, in *Conférence Moshé Flato 1999, Vol. II (Dijon)*, volume 22 of *Math. Phys. Stud.*, pages 307–331, Kluwer Acad. Publ., Dordrecht, 2000.